

# Solution to Atiyah and MacDonald

## Chapter 2. Modules

Jaehyeon Lee

Last update: April 22, 2024

This is a solution to Exercise problems in Chapter 2 of "Introduction to Commutative Algebra" written by M. F. Atiyah and I. G. MacDonald. You can find the updated version and solutions to other chapters on my personal website: [\[https://ijhlee0511.github.io\]](https://ijhlee0511.github.io).

**WARNING** This solution is written for self-study purposes and to consolidate my understanding. **I do not take responsibility for any disadvantages resulting from the use of this solution. It is at your own risk.** If you find any typos or errors in this solution, please feel free to contact me via email at [\[ijhlee0511@gmail.com\]](mailto:ijhlee0511@gmail.com) or [\[ijhlee0511@kaist.ac.kr\]](mailto:ijhlee0511@kaist.ac.kr).

### Exercises and Solutions

**2.1.** Show that  $(\mathbb{Z}/m\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/n\mathbb{Z}) = 0$  if  $m, n$  are coprime.

**Solution.** There exist integers  $x$  and  $y$  such that  $xm + yn = 1$ . As a result, for any  $a \in \mathbb{Z}/m\mathbb{Z}$  and  $b \in \mathbb{Z}/n\mathbb{Z}$ ,

$$a \otimes b = (xm + yn)(a \otimes b) = xma \otimes b + a \otimes ynb = 0.$$

□

**2.2.** Let  $A$  be a ring,  $\alpha$  an ideal,  $M$  an  $A$ -module. Show that  $(A/\alpha) \otimes_A M$  is isomorphic to  $M/\alpha M$ .

**Solution.** By the natural inclusion and projection, a sequence  $0 \rightarrow \alpha \rightarrow A \rightarrow A/\alpha \rightarrow 0$  is exact. Tensoring the sequence, we get an exact sequence  $\alpha \otimes M \rightarrow M \rightarrow A/\alpha \otimes M \rightarrow 0$ . Since the map  $\alpha \otimes M \rightarrow M$  in the sequence is given by  $x \otimes m \mapsto xm$ , we get  $A/\alpha \otimes M \cong M/\text{Im}(\alpha \otimes M \rightarrow M) \cong M/\alpha M$ . □

**2.3.** Let  $A$  be a local ring,  $M$  and  $N$  finitely generated  $A$ -modules. Prove that if  $M \otimes N = 0$ , then  $M = 0$  or  $N = 0$ .

**Solution.** Let  $\mathfrak{m}$  be the unique maximal ideal. Suppose neither  $M = 0$  nor  $N = 0$ . Then by Nakayama's lemma,  $M/\mathfrak{m}M \neq 0$  and  $N/\mathfrak{m}N \neq 0$ . We can view  $M/\mathfrak{m}M$  and  $N/\mathfrak{m}N$  as finite dimensional vector spaces over  $k := A/\mathfrak{m}$ . Therefore,  $V := M/\mathfrak{m}M \otimes_k N/\mathfrak{m}N$  is a  $k$ -vector space with dimension  $(\dim_k M/\mathfrak{m}M) \times (\dim_k N/\mathfrak{m}N)^1$ , implying it is nonzero.

---

<sup>1</sup>There are a lot of ways to show this, but one may use the fact that every  $n$ -dimensional vector space is isomorphic to  $k^n$ . In general,  $R^n \otimes_R R^m = R^{nm}$  for any commutative ring  $R$  by Proposition 2.14.

Define a map  $f : M/\mathfrak{m}M \times N/\mathfrak{m}N \rightarrow V$  by  $(\bar{m}, \bar{n}) \mapsto \bar{m} \otimes_k \bar{n}$ . Viewing  $V$  as an  $A$ -module, this is naturally an  $A$ -bilinear map, so there is a surjective  $A$ -module homomorphism  $f^* : M/\mathfrak{m}M \otimes_A N/\mathfrak{m}N \twoheadrightarrow V$  sending  $\bar{m} \otimes_A \bar{n}$  to  $\bar{m} \otimes_k \bar{n}$ . From the natural surjection  $N \twoheadrightarrow N/\mathfrak{m}N$  we get an exact sequence  $M/\mathfrak{m}M \otimes_A N \rightarrow M/\mathfrak{m}M \otimes_A N/\mathfrak{m}N \rightarrow 0$ ; hence, there is a surjective  $A$ -module homomorphism  $M/\mathfrak{m}M \otimes_A N \twoheadrightarrow V$ . However, by Exercise 2.2,

$$M/\mathfrak{m}M \otimes_A N = (A/\mathfrak{m} \otimes_A M) \otimes_A N = A/\mathfrak{m} \otimes_A (M \otimes_A N) = 0,$$

a contradiction.  $\square$

**2.4.** Let  $M_i$  ( $i \in I$ ) be any family of  $A$ -modules, and let  $M$  be their direct sum. Prove that  $M$  is flat  $\Leftrightarrow$  each  $M_i$  is flat.

**Solution.** Suppose  $M$  is flat. If  $f : N' \rightarrow N$  is an injective  $A$ -module homomorphism, then  $f \otimes 1 : N \otimes M \rightarrow N' \otimes M$ . However,  $N \otimes M = \bigoplus_{i \in I} N \otimes M_i$  and  $N' \otimes M = \bigoplus_{i \in I} N' \otimes M_i$ . Observing  $(f \otimes 1)(N \otimes M_i) \subseteq N' \otimes M_i$ , the restriction  $(f \otimes 1)|_{N \otimes M_i} : N \otimes M_i \rightarrow N' \otimes M_i$  is also injective. Therefore, each  $M_i$  is also flat.

Conversely, suppose each  $M_i$  is flat. If  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  is an exact sequence of  $A$ -modules, then  $0 \rightarrow N' \otimes M_i \rightarrow N \otimes M_i \rightarrow N'' \otimes M_i \rightarrow 0$  is also exact. Therefore, the direct sum of exact sequences  $0 \rightarrow \bigoplus_{i \in I} N' \otimes M_i \rightarrow \bigoplus_{i \in I} N \otimes M_i \rightarrow \bigoplus_{i \in I} N'' \otimes M_i \rightarrow 0$  is also exact.  $\square$

**2.5.** Let  $A[x]$  be the ring of polynomials in one indeterminate over a ring  $A$ . Prove that  $A[x]$  is a flat  $A$ -algebra.

**Solution.** As an  $A$ -module,  $A[x] = \bigoplus_{i=0}^{\infty} Ax^i$  where  $Ax^i \cong A$  for each  $i \in \mathbb{Z}_{\geq 0}$ . Since  $A$  is clearly flat, its direct sum  $A[x] \cong A^{\mathbb{Z}_{\geq 0}}$  is also flat by Exercise 2.4.  $\square$

**2.6.** For any  $A$ -module, let  $M[x]$  denote the set of all polynomials in  $x$  with coefficients in  $M$ , that is to say expressions of the form

$$m_0 + m_1x + \cdots + m_rx^r \quad (m_i \in M).$$

Defining the product of an element of  $A[x]$  and an element of  $M[x]$  in the obvious way, show that  $M[x]$  is an  $A[x]$ -module.

Show that  $M[x] \cong A[x] \otimes_A M$ .

**Solution.**  $M[x]$  is clearly an abelian group since  $M$  is itself an abelian group. Precisely, its abelian group structure is the same with the direct sum  $M[x] \cong \bigoplus_{i=1}^{\infty} Mx^i \cong \bigoplus_{i=1}^{\infty} M$ . Also, rules for scalar multiplication by  $A[x]$  straightforwardly (but too tedious to type every minor detail) hold.

Firstly, let's construct an  $A$ -module isomorphism  $A[x] \otimes_A M \rightarrow M[x]$ , and show that it preserves  $A[x]$ -scalar multiplication later. Define a map  $f : A[x] \times M \rightarrow M[x]$  by

$$(a_0 + a_1x + \cdots + a_rx^r, m) \mapsto a_0m + (a_1m)x + \cdots + (a_rm)x^r.$$

It is easy to verify  $f$  is  $A$ -bilinear. Therefore, there is a unique  $A$ -module homomorphism  $\phi : A[x] \otimes_A M \rightarrow M[x]$  sending  $a_ix^i \otimes m$  to  $(a_im)x^i$ . Now, define an  $A$ -module homomorphism  $\psi : M[x] \rightarrow A[x] \otimes_A M$  by  $m_ix^i \mapsto x^i \otimes m_i$  for each  $i \in \mathbb{Z}_{\geq 0}$ .<sup>2</sup> Then  $\psi \circ \phi = \text{id}_{A[x] \otimes M}$  and

<sup>2</sup>Since  $M[x] \cong \bigoplus_i Mx^i$  as an  $A$ -module, this assignment uniquely determines  $\psi$ , which is well-defined.

$\phi \circ \psi = \text{id}_{M[x]}$ ,<sup>3</sup> so  $\phi : A[x] \otimes_A M \rightarrow M[x]$  is an  $A$ -module isomorphism. We claim that  $\phi$  actually respects scalar multiplication by  $A[x]$ . For  $b_0 + b_1x + \cdots + b_sx^s \in A$ , we have

$$\begin{aligned} f((b_0 + b_1x + \cdots + b_sx^s)(a_ix^i \otimes m)) &= f((a_ib_0x^i + a_ib_1x^{i+1} + \cdots + a_ib_sx^{s+i}) \otimes m) \\ &= a_ib_0mx^i + a_ib_1mx^{i+1} + \cdots + a_ib_smx^{s+i} \\ &= (b_0 + b_1x + \cdots + b_sx^s)(a_imx^i) \\ &= (b_0 + b_1x + \cdots + b_sx^s)f(a_ix^i \otimes m), \end{aligned}$$

for each  $i \in \mathbf{Z}_{\geq 0}$ . Therefore,  $\phi$  is also an  $A[x]$ -module isomorphism.  $\square$

**2.7.** Let  $\mathfrak{p}$  be a prime ideal in  $A$ . Show that  $\mathfrak{p}[x]$  is a prime ideal in  $A[x]$ . If  $\mathfrak{m}$  is a maximal ideal in  $A$ , is  $\mathfrak{m}[x]$  a maximal ideal in  $A[x]$ ?

**Solution.** Since  $\mathfrak{p}[x]$  is the kernel of  $A[x] \rightarrow (A/\mathfrak{p})[x]$ , we have  $A[x]/\mathfrak{p}[x] \cong (A/\mathfrak{p})[x]$ . Suppose there are  $a_0 + \cdots + a_rx^r$  and  $b_0 + \cdots + b_sx^s$  in  $(A/\mathfrak{p})[x]$  with non-zero  $a_r$  and  $b_s$  so that  $(a_0 + \cdots + a_rx^r)(b_0 + \cdots + b_sx^s) = 0$ . However, we get  $a_rb_s = 0$ , a contradiction for  $A/\mathfrak{p}$  is an integral domain. As a result,  $\mathfrak{p}[x]$  is a prime ideal in  $A[x]$ .

Let  $k$  be a field,  $A = k[y]$ , and  $\mathfrak{m} = (y)$ . Then  $k[y][x]/(y)[x] \cong (k[y]/(y))[x] \cong k[x]$ , which is clearly not a field in general. Therefore,  $\mathfrak{m}[x]$  is not a maximal ideal.  $\square$

**2.8.** i) If  $M$  and  $N$  are flat  $A$ -modules, then so is  $M \otimes_A N$ .

ii) If  $B$  is a flat  $A$ -algebra and  $N$  is a flat  $B$ -module, then  $N$  is flat as an  $A$ -module.

**Solution.** i) If  $f : L' \rightarrow L$  is an injective  $A$ -module homomorphism, then  $(1 \otimes f) : N \otimes L' \rightarrow N \otimes L$  is injective. Also,  $1 \otimes (1 \otimes f) : M \otimes (N \otimes L') \rightarrow M \otimes (N \otimes L)$  is injective. By the identification  $M \otimes (N \otimes L') = (M \otimes N) \otimes L'$  and  $M \otimes (N \otimes L) = (M \otimes N) \otimes L$ , it induces an injective map  $1' \otimes f : (M \otimes N) \otimes L' \rightarrow (M \otimes N) \otimes L$  given by  $(m \otimes n) \otimes l' \mapsto (m \otimes n) \otimes l$ . This shows  $M \otimes_A N$  is flat.

ii) If  $f : M' \rightarrow M$  is an injective  $A$ -module homomorphism then  $(1 \otimes f) : B \otimes_A M' \rightarrow B \otimes_A M$  is injective. However, we can regard this injective map as a  $B$ -module homomorphism, since  $b_1b_2 \otimes m' \mapsto b_1b_2 \otimes f(m') = b_1(b_2 \otimes f(m'))$ . Therefore, we get an injective  $B$ -module homomorphism  $N \otimes_B (B \otimes_A M') \rightarrow N \otimes_B (B \otimes_A M)$ , but by the canonical isomorphism in 2.15 of the main text, we get  $N \otimes_A M' \rightarrow N \otimes_A M$  given by  $n \otimes m' \mapsto n \otimes f(m')$ . As a result,  $N$  is flat as an  $A$ -module.  $\square$

**2.9.** Let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence of  $A$ -modules. If  $M'$  and  $M''$  are finitely generated, then so is  $M$ .

**Solution.** Regard  $M'$  as a submodule of  $M$ . Since  $M/M' \cong M''$ , there are  $x_1, \dots, x_n \in M$  so that  $x_1 + M', \dots, x_n + M'$  generate  $M/M'$ . That is, every element of  $M$  belongs to some coset, which is a linear combination of  $x_1 + M', \dots, x_n + M'$ . Let  $y_1, \dots, y_m$  be generators of  $M'$ . Then  $x_1, \dots, x_n, y_1, \dots, y_m$  generate  $M$ .  $\square$

**2.10.** Let  $A$  be a ring,  $\alpha$  an ideal contained in the Jacobson radical of  $A$ ; let  $M$  be an  $A$ -module and  $N$  a finitely generated  $A$ -module, and let  $u : M \rightarrow N$  be a homomorphism. If the induced homomorphism  $M/\alpha M \rightarrow N/\alpha N$  is surjective, then  $u$  is surjective.

<sup>3</sup>Checking only for generators suffices to show this.

**Solution.** Since  $N$  is finitely generated, so is  $N/\alpha N$ . Suppose  $n_1 + \alpha N, \dots, n_r + \alpha N$  generate  $N/\alpha N$  where  $n_i = u(m_i)$  for some  $m_i \in M$ . Then  $N = \alpha N + \sum_{i=1}^r \alpha N n_i$ , so

$$N / (\sum_{i=1}^r \alpha N n_i) \cong \alpha N / (\alpha N \cap \sum_{i=1}^r \alpha N n_i) \cong \alpha (N / \sum_{i=1}^r \alpha N n_i).$$

By Nakayama's lemma, we get  $N = \sum_{i=1}^r \alpha N n_i$ . Therefore,  $u$  is surjective.  $\square$

**2.11.** Let  $A$  be a ring  $\neq 0$ . Show that  $A^m \cong A^n \Rightarrow m = n$ .

If  $\phi : A^m \rightarrow A^n$  is surjective, then  $m \geq n$ .

If  $\phi : A^m \rightarrow A^n$  is injective, is it always the case  $m \leq n$ ?

**Solution.** Let  $\mathfrak{m}$  be a maximal ideal of  $A$  and let  $\phi : A^m \rightarrow A^n$  be an isomorphism. Then  $1 \otimes \phi : (A/\mathfrak{m}) \otimes A^m \rightarrow (A/\mathfrak{m}) \otimes A^n$  is an isomorphism between vector spaces of dimensions  $m$  and  $n$  over the field  $k = A/\mathfrak{m}$ , since  $(A/\mathfrak{m}) \otimes A^m \cong k^m$  and  $(A/\mathfrak{m}) \otimes A^n \cong k^n$  by Exercise 2.2. Hence,  $m = n$ .

If  $\phi$  is surjective, then the tensored morphism  $1 \otimes \phi : k^m \rightarrow k^n$  is again surjective by Proposition 2.18. Therefore,  $m \geq n$  by linear algebra.

The injectivity part is a very famous problem, and there are a lot of good answers for it. One way with a structural approach involves exterior algebra as Corollary 5.11 of [1]. However, I can not figure out a better answer than the following solution [2] in MathOverflow, which uses only Proposition 2.4. Suppose there is an injective  $A$ -module homomorphism  $\phi : A^m \rightarrow A^n$  with  $m > n$ . Identifying  $A^n$  with  $\{(a_1, \dots, a_n, 0, \dots, 0) \in A^m \mid a_i \in A\} \subseteq A^m$ , we can regard it as an  $A$ -module embedding  $\phi : A^m \hookrightarrow A^m$ ; i.e.,  $\phi(A^m) \subseteq A^m$ . Therefore, by Proposition 2.4,  $\phi$  satisfies an equation of the form

$$p(\phi) = \phi^d + a_1 \phi^{d-1} + \dots + a_d = 0$$

where  $a_i$  are in  $A$  and  $p(x) \in A[x]$ . Suppose the polynomial  $p$  has the minimum degree (well ordering principle). If  $a_d = 0$ , then  $\phi(\phi^{d-1} + a_1 \phi^{d-2} + \dots + a_{d-1})(v) = 0$  for all  $v \in A^m$ . However, by the injectivity of  $\phi$ , we get  $\phi^{d-1} + a_1 \phi^{d-2} + \dots + a_{d-1} = 0$ , a contradiction. Therefore  $a_d$  is nonzero. However, the  $m$ -th coordinate of  $p(\phi)(0, \dots, 0, 1)$  is  $a_d$ , contradicting to the assumption  $\phi(A^m) \subseteq A^n$ . This shows  $m \leq n$ .  $\square$

**2.12.** Let  $M$  be a finitely generated  $A$ -module and  $\phi : M \rightarrow A^n$  a surjective homomorphism. Show that  $\text{Ker}(\phi)$  is finitely generated.

**Solution.** Let  $e_1, \dots, e_n$  be a basis of  $A^n$  and choose  $u_i \in M$  such that  $\phi(u_i) = e_i$  ( $1 \leq i \leq n$ ). Let  $N$  be the submodule of  $M$  generated by  $u_1, \dots, u_n$ . Then every element of  $x$  of  $M$  must be in some coset  $y + \text{Ker}(\phi)$  for some  $y \in N$  if and only if  $\phi(x) = \phi(y)$ . This shows  $N + \text{Ker}(\phi) = M$ . If  $r_1 u_1 + \dots + r_n u_n \in N$  is in  $\text{Ker}(\phi)$ , then  $r_1 e_1 + \dots + r_n e_n = 0$  in  $A^n$ . This implies  $r_1 = \dots = r_n = 0$ , so  $N \cap \text{Ker}(\phi) = 0$ . As a result,  $M = N \oplus \text{Ker}(\phi)$ . Since  $M$  is finitely generated, its quotient  $M/N \cong \text{Ker}(\phi)$  is also finitely generated.  $\square$

**2.13.** Let  $f : A \rightarrow B$  be a ring homomorphism, and let  $N$  be a  $B$ -module. Regarding  $N$  as an  $A$ -module by restriction of scalars, form the  $B$ -module  $N_B = B \otimes_A N$ . Show that the homomorphism  $g : N \rightarrow N_B$  which maps  $y$  to  $1 \otimes y$  is injective and that  $g(N)$  is a direct summand of  $N_B$ .

**Solution.** We shall show  $1 \otimes y \in B \otimes_A N$  is zero only if  $y = 0$ . Define  $p : N_B \rightarrow N$  by  $p(b \otimes y) = by$ . It is easy to see that  $p$  is a  $B$ -module homomorphism. However,  $p(1 \otimes y) = 0$  only if  $y = 0$ , so  $g$  is injective. Since  $p$  is clearly surjective, we have  $N_B / \text{Ker}(p) \cong N \cong \text{Im}(g)$ . Let  $p^* : N_B / \text{Ker}(p) \rightarrow N$  be the induced map by  $p$ . Then  $p^*(1 \otimes y + \text{Ker}(p)) = y$  for all  $y \in N$ , so  $\text{Im}(g) + \text{Ker}(p) = N_B$  since every element of  $N$  must be in some coset  $1 \otimes y + \text{Ker}(p)$ . However, because  $p \circ g = \text{id}_N$ , we get  $\text{Im}(g) \cap \text{Ker}(p) = 0$ . This shows  $N_B \cong \text{Im}(g) \oplus \text{Ker}(p) \cong N \oplus \text{Ker}(p)$ .  $\square$

**2.14.** A partially ordered set  $I$  is said to be a *directed set* if for each pair  $i, j$  in  $I$  there exists  $k \in I$  such that  $i \leq k$  and  $j \leq k$ .

Let  $A$  be a ring, let  $I$  be a directed set and let  $(M_i)_{i \in I}$  be a family of  $A$ -modules indexed by  $I$ . For each pair  $i, j$  in  $I$  such that  $i \leq j$ , let  $\mu_{ij} : M_i \rightarrow M_j$  be an  $A$ -homomorphism, and suppose that the following axioms are satisfied:

- (1)  $\mu_{ii}$  is the identity mapping of  $M_i$ , for all  $i \in I$ ;
- (2)  $\mu_{ik} = \mu_{jk} \circ \mu_{ij}$  whenever  $i \leq j \leq k$ .

Then the modules  $M_i$  and homomorphisms  $\mu_{ij}$  are said to form a *direct system*  $\mathbf{M} = (M_i, \mu_{ij})$  over the directed set  $I$ .

We shall construct an  $A$ -module  $M$  called the *direct limit* of the direct system  $\mathbf{M}$ . Let  $C$  be the direct sum of the  $M_i$ , and identify each module  $M_i$  with its canonical image in  $C$ . Let  $D$  be the submodule of  $C$  generated by all elements of the form  $x_i - \mu_{ij}(x_i)$  where  $i \leq j$  and  $x_i \in M_i$ . Let  $M = C/D$ , let  $\mu : C \rightarrow M$  be the projection and let  $\mu_i$  be the restriction of  $\mu$  to  $M_i$ .

The module  $M$ , or more correctly the pair consisting of  $M$  and the family of homomorphisms  $\mu_i : M_i \rightarrow M$ , is called the *direct limit* of the direct system  $\mathbf{M}$ , and is written  $\varinjlim M_i$ . From the construction it is clear that  $\mu_i = \mu_j \circ \mu_{ij}$  whenever  $i \leq j$ .

**Solution.** There is nothing to do.  $\square$

**2.15.** In the situation of Exercise 14, show that every element of  $M$  can be written in the form  $\mu_i(x_i)$  for some  $i \in I$  and some  $x_i \in M_i$ .

Show also that if  $\mu_i(x_i) = 0$  then there exists  $j \geq i$  such that  $\mu_{ij}(x_i) = 0$  in  $M_j$ .

**Solution.** In the construction of  $M$  in Exercise 2.14, there are finitely many  $i_1, \dots, i_n \in I$  so that  $y = \mu_{i_1}(x_{i_1}) + \dots + \mu_{i_n}(x_{i_n})$ . Choose some  $j \geq i_1, \dots, i_n$  and let  $x_j := \mu_{i_1 j}(x_{i_1}) + \dots + \mu_{i_n j}(x_{i_n})$ . Then  $\mu_j(x_j) = y$ .

Suppose  $\mu_i(x_i) = 0$ . Then  $x_i$  is in  $\text{Ker}(\mu)$ , where  $\mu : C := \bigoplus_{i \in I} M_i \rightarrow M$  is the projection. Since  $\text{Ker}(\mu)$  is a submodule of  $C$  generated by  $\{y_i - \mu_{ij}(y_i) \mid i, j \in I, j \geq i\}$ , we have  $x_i = \sum_{p=1}^n (y_p - \mu_{i_p j_p}(y_p))$  where  $y_p \in M_{i_p}$  and  $i_p \leq j_p$ . However, every term not in  $M_i$  must be eliminated by other terms, so  $x_i = y_{i'} - \mu_{i' j}(y_{i'})$  for some  $y_{i'} \in M_{i'}$ . The only possible way is  $y_{i'} = x_i$  and  $\mu_{ij}(x_i) = 0$ .  $\square$

**2.16.** Show that the direct limit is characterized (up to isomorphism) by the following property. Let  $N$  be an  $A$ -module and for each  $i \in I$  let  $\alpha_i : M_i \rightarrow N$  be an  $A$ -module homomorphism such that  $\alpha_i = \alpha_j \circ \mu_{ij}$  whenever  $i \leq j$ . Then there exists a unique homomorphism  $\alpha : M \rightarrow N$  such that  $\alpha_i = \alpha \circ \mu_i$  for all  $i \in I$ .

**Solution.** Rephrasing the property as a commutative diagram is as follows:

$$\begin{array}{ccc}
 M_i & \xrightarrow{\mu_{ij}} & M_j \\
 \searrow \mu_i & & \swarrow \mu_j \\
 & M & \\
 \swarrow \alpha_i & \downarrow \exists! \alpha & \searrow \alpha_j \\
 & N &
 \end{array}$$

for  $j \geq i$ , where each triangle in the diagram commutes. This property uniquely characterizes  $M$  up to ‘unique’ isomorphism. Suppose our given  $(M, \mu_i)$  satisfies the property, and another module  $(M', \mu'_i)$  satisfies the same property. Trivially, if we plug  $M$  to  $N$ , the unique morphism  $\alpha$  is the identity  $\text{id}_M$ . However, by the assumption, there are unique morphisms  $\beta$  and  $\gamma$  so that each triangle of the following diagram commutes

$$\begin{array}{ccccc}
 M_i & & & & \\
 \downarrow \mu_{ij} & \searrow \mu_i & & \searrow \mu_j & \\
 & M & \xrightarrow{\exists! \beta} & M' & \xrightarrow{\exists! \gamma} M \\
 & \swarrow \mu_j & & \swarrow \mu'_j & \\
 M_j & & & &
 \end{array}$$

for each  $j \geq i$ . Therefore we get  $\gamma \circ \beta = \text{id}_M$ . Ditto  $\beta \circ \gamma = \text{id}_{M'}$ , and this shows  $M$  and  $M'$  are isomorphic by ‘unique’ isomorphisms  $\beta$  and  $\gamma$ .

Now let’s show  $(M, \mu_i)$  actually satisfy the property. Define an  $A$ -module homomorphism  $f : \bigoplus_{i \in I} M_i \rightarrow N$  as  $(x_i)_{i \in I} \mapsto \sum_{i \in I} \alpha_i(x_i)$ . Since  $\alpha_i(x_i) = \alpha_j(\mu_{ij}(x_i))$  by the assumption,  $f(x_i - \mu_{ij}(x_i)) = 0$ . Therefore  $\text{Ker}(\mu) \subseteq \text{Ker}(f)$ , so we get the induced  $A$ -module homomorphism  $\alpha : M \rightarrow N$  satisfying  $\alpha \circ \mu = f$ . By the construction,  $\alpha(\mu_i(x_i)) = f(x_i) = \alpha_i(x_i)$  for any  $x_i \in M_i$ , so  $\alpha$  satisfies the desired property. To show the uniqueness of  $\alpha$ , suppose  $\alpha' : M \rightarrow N$  also satisfies the same property. By Exercise 2.15, every element of  $M$  can be written in the form  $\mu_i(x_i)$  for some  $x_i \in M_i$ . But  $\alpha'(\mu_i(x_i)) = \alpha_i(x_i) = \alpha(\mu_i(x_i))$ . This ends the proof.  $\square$

**2.17.** Let  $(M_i)_{i \in I}$  be a family of submodules of an  $A$ -module, such that for each pair of indices  $i, j$  in  $I$  there exists  $k \in I$  such that  $M_i + M_j \subseteq M_k$ . Define  $i \leq j$  to mean  $M_i \subseteq M_j$  and let  $\mu_{ij} : M_i \rightarrow M_j$  be the embedding of  $M_i$  in  $M_j$ . Show that

$$\varinjlim M_i = \sum M_i = \bigcup M_i.$$

In particular, any  $A$ -module is the direct limit of its finitely generated submodules.

**Solution.** For any  $m, n \in \sum M_i$ , notice  $m + n$  belongs to some ambient module  $M_k$ , so  $\bigcup M_i = \sum M_i$ . Let  $\mu_i : M_i \rightarrow \sum M_i$  be the natural inclusion, and suppose  $N$  be an  $A$ -module and for each  $i \in I$  let  $\alpha_i : M_i \rightarrow N$  is an  $A$ -module homomorphism such that  $\alpha_i = \alpha_j \circ \mu_{ij}$  whenever  $i \leq j$ . Then an  $A$ -module homomorphism  $\alpha : \sum M_i \rightarrow N$  given by  $(x_i)_{i \in I} \mapsto \sum \alpha_i(x_i)$  satisfies  $\alpha(\mu_i(x_i)) = \alpha_i(x_i)$  for arbitrary  $x_i \in M_i$ . Since  $\mu_i$  is nothing but inclusion, such  $\alpha$  satisfying  $\alpha = \alpha_i \circ \mu_i$  for any  $i \in I$  is unique. This shows that  $\varinjlim M_i = \sum M_i$  by Exercise 2.16.

In particular, let  $M$  be any  $A$ -module, and  $(M_i)_{i \in I}$  be the collection of all finitely generated submodules of  $M$ . For each  $i, j \in I$ ,  $M_i + M_j$  is also finitely generated; hence  $I$  and  $(M_i)_{i \in I}$  satisfies the desired property. Moreover, for any  $x \in M$ ,  $Ax$  is a finitely generated submodule itself, so  $M = \sum M_i = \varinjlim M_i$ .  $\square$

**2.18.** Let  $\mathbf{M} = (M_i, \mu_{ij})$ ,  $\mathbf{N} = (N_i, v_{ij})$  be direct systems of  $A$ -modules over the same directed set. Let  $M, N$  be the direct limits and  $\mu_i : M_i \rightarrow M$ ,  $v_i : N_i \rightarrow N$  the associated homomorphisms.

A homomorphism  $\Phi : \mathbf{M} \rightarrow \mathbf{N}$  is by definition a family of  $A$ -module homomorphisms  $\phi_i : M_i \rightarrow N_i$ , such that  $\phi_j \circ \mu_{ij} = v_{ij} \circ \phi_i$  whenever  $i \leq j$ . Show that  $\Phi$  defines a unique homomorphism  $\phi = \varinjlim \phi_i : M \rightarrow N$  such that  $\phi \circ \mu_i = v_i \circ \phi_i$  for all  $i \in I$ .

**Solution.** Let  $\alpha_i := v_i \circ \phi_i$  for each  $i \in I$ . Then by the assumption, we get

$$\begin{aligned} \alpha_i &= v_i \circ \phi_i \\ &= v_j \circ v_{ij} \circ \phi_i \\ &= v_j \circ \phi_j \circ \mu_{ij} \\ &= \alpha_j \circ \mu_{ij}, \end{aligned}$$

whenever  $i \leq j$ . By Exercise 2.16, this implies that there exists a unique homomorphism  $\phi : M \rightarrow N$  so that the following diagram commutes:

$$\begin{array}{ccccc} M_j & \xrightarrow{\phi_j} & N_j & & \\ & \searrow \mu_j & \swarrow v_j & & \\ & & M & \xrightarrow{\exists! \phi} & N \\ & \nearrow \mu_i & \nwarrow v_i & & \\ M_i & \xrightarrow{\phi_i} & N_i & & \end{array}$$

whenever  $i \leq j$ . This ends the proof.  $\square$

**2.19.** A sequence of direct systems and homomorphisms

$$\mathbf{M} \rightarrow \mathbf{N} \rightarrow \mathbf{P}$$

is *exact* if the corresponding sequence of modules and module homomorphisms is exact for each  $i \in I$ . Show that the sequence  $M \rightarrow N \rightarrow P$  of direct limits is then exact.

**Solution.** For the notations, let  $\mathbf{M} = (M_i, \mu_{ij})$ ,  $\mathbf{N} = (N_i, v_{ij})$ , and  $\mathbf{P} = (P_i, \pi_{ij})$  be direct systems of  $A$ -modules over the same directed set  $I$ . Let  $M, N$ , and  $P$  be the direct limits and  $\mu_i : M_i \rightarrow M$ ,  $v_i : N_i \rightarrow N$ , and  $\pi_i : P_i \rightarrow P$  be the associated homomorphisms. Let  $\Phi : \mathbf{M} \rightarrow \mathbf{N}$  and  $\Psi : \mathbf{N} \rightarrow \mathbf{P}$  denote homomorphisms of direct systems so that the given sequence is exact where  $\phi_i : M_i \rightarrow N_i$ , and  $\psi_i : N_i \rightarrow P_i$  are associated homomorphisms. By Exercise 2.18, they define a unique homomorphism  $\phi = \varinjlim \phi_i : M \rightarrow N$  and  $\psi = \varinjlim \psi_i : N \rightarrow P$  such that  $\phi \circ \mu_i = v_i \circ \phi_i$  and  $\psi \circ v_i = \pi_i \circ \psi_i$ . Then the following diagram commutes:

$$\begin{array}{ccccccc} M_i & \xrightarrow{\phi_i} & N_i & \xrightarrow{\psi_i} & P_i & & \\ \downarrow \mu_i & \searrow \mu_{ij} & \downarrow v_i & \searrow v_{ij} & \downarrow \pi_i & \searrow \pi_{ij} & \\ & & M_j & \xrightarrow{\phi_j} & N_j & \xrightarrow{\psi_j} & P_j \\ & \nearrow \mu_j & \downarrow v_j & \nearrow v_j & \downarrow \pi_j & \nearrow \pi_j & \\ M & \xrightarrow{\phi} & N & \xrightarrow{\psi} & P & & \end{array}$$

whenever  $i \leq j$ , where the first and second rows are exact by the assumption. We claim that  $\text{Im}(\phi) = \text{Ker}(\psi)$ ; i.e., the third row is exact. By Exercise 2.15, for any  $m \in M$ , there are some  $i \in I$  and some  $m_i \in M_i$  so that  $m = \mu_i(m_i)$ . Then  $\psi(\phi(m)) = (\psi \circ \phi \circ \mu_i)(m_i) = (\pi_i \circ \psi_i \circ \phi_i)(m_i) = \pi_i(0) = 0$ , so  $\text{Im}(\phi) \subseteq \text{Ker}(\psi)$ . For the reverse inclusion, suppose  $n$  is in  $\text{Ker}(\psi)$ . By Exercise 2.15 again, there exists some  $i \in I$  and some  $n_i \in N_i$  such that  $n = v_i(n_i)$ . However,  $0 = \psi(n) = \psi(v_i(n_i)) = \pi_i(\psi_i(n_i))$ , so there exists some  $j \geq i$  such that  $\pi_{ij}(\psi_i(n_i)) = 0$  by the second statement of Exercise 2.15. Notice  $\psi_j(v_{ij}(n_i)) = \pi_{ij}(\psi_i(n_i)) = 0$ . Thus, there exists some  $m_j \in M_j$  such that  $\phi_j(m_j) = v_{ij}(n_i)$  due to the assumption that  $\text{Im}(\phi_j) = \text{Ker}(\psi_j)$ . As a result,

$$\phi(\mu_j(m_j)) = v_j(\phi_j(m_j)) = v_j(v_{ij}(n_i)) = v_i(n_i) = n.$$

This shows  $\text{Im}(\phi) = \text{Ker}(\psi)$ . □

**2.20.** Keeping the same notation as in Exercise 14, let  $N$  be any  $A$ -module. Then  $(M_i \otimes N, \mu_{ij} \otimes 1)$  is a direct system; let  $P = \varinjlim (M_i \otimes N)$  be its direct limit. For each  $i \in I$  we have a homomorphism  $\mu_i \otimes 1 : M_i \otimes N \rightarrow M \otimes N$ , hence by Exercise 16 a homomorphism  $\psi : P \rightarrow M \otimes N$ . Show that  $\psi$  is an isomorphism, so that

$$\varinjlim (M_i \otimes N) \cong (\varinjlim M_i) \otimes N.$$

**Solution.** For the notation, let  $\mu'_i : M_i \otimes N \rightarrow P$  denote the canonical  $A$ -module homomorphism characterizing the direct limit  $P$ . Then  $\psi \circ \mu'_i = \mu_i \otimes 1$  for all  $i \in I$ . For each  $i \in I$ , let  $g_i : M_i \times N \rightarrow M_i \otimes N$  be the canonical bilinear mapping given by  $(m_i, n) \mapsto m_i \otimes n$ . Fixing  $n \in N$ , we get an  $A$ -module homomorphisms  $g_i(-, n) : M_i \rightarrow M_i \otimes N$ , and it is easy to see that they form a homomorphism  $(M_i, \mu_{ij}) \rightarrow (M_i \otimes N, \mu_{ij} \otimes 1)$  between two directed system. Therefore, by Exercise 2.18, they define a unique homomorphism  $g(-, n) : M \rightarrow P$  such that  $g(-, n) \circ \mu_i = \mu'_i \circ g_i(-, n)$ . We claim that  $g(m, -) : N \rightarrow P$  is also an  $A$ -module homomorphism for each fixed  $m \in M$ . By Exercise 2.15, there exist some  $i \in I$  and some  $m_i \in M_i$  so that  $\mu_i(m_i) = m$ . Then for any  $n_1, n_2 \in N$  and  $a \in A$ , we have

$$\begin{aligned} g(m, n_1 + an_2) &= g(\mu_i(m_i), n_1 + an_2) \\ &= (g(-, n_1 + an_2) \circ \mu_i)(m_i) \\ &= \mu'_i(g_i(m_i, n_1 + an_2)) \\ &= \mu'_i(g_i(m_i, n_1) + ag_i(m_i, n_2)) \\ &= \mu'_i(g_i(m_i, n_1)) + a\mu'_i(g_i(m_i, n_2)) \\ &= g(m, n_1) + ag(m, n_2), \end{aligned}$$

assuming  $m = \mu_i(m_i)$  for some  $i \in I$  and some  $m_i \in M_i$ . We finally get a bilinear map  $g : M \times N \rightarrow P$ , and hence we obtain the corresponding  $A$ -module homomorphism  $\phi : M \otimes N \rightarrow P$  such that  $\phi(m \otimes n) = g(m, n)$ .

Now we claim that  $\phi$  and  $\psi$  are mutually inverse. For any  $m \in M$  and  $n \in N$ , assuming



$m = \mu_i(m_i)$  for some  $i \in I$  and some  $m_i \in M_i$ ,

$$\begin{aligned}
 \psi(\phi(m \otimes n)) &= \psi(g(m, n)) \\
 &= \psi(g(\mu_i(m_i), n)) \\
 &= (\psi \circ \mu'_i)(g_i(m_i, n)) \\
 &= (\mu_i \otimes 1)(g_i(m_i, n)) \\
 &= (\mu_i \otimes 1)(m_i \otimes n) \\
 &= \mu_i(m_i) \otimes n \\
 &= m \otimes n,
 \end{aligned}$$

so  $\psi \circ \phi = \text{id}_{M \otimes N}$ . For the converse, for given  $p \in P$ , there is some  $i \in I$  and some  $x_i \in M_i \otimes N$  so that  $p = \mu'_i(x_i)$ . We may write  $x_i = \sum_{j=1}^k m_{ij} \otimes n_j$  for some  $k \in \mathbf{Z}_{\geq 0}$ ,  $m_{i1}, \dots, m_{ik} \in M_i$ , and  $n_1, \dots, n_k \in N$ . Then,

$$\begin{aligned}
 \phi(\psi(p)) &= (\phi \circ \psi \circ \mu'_i)(x_i) \\
 &= \sum_{j=1}^k (\phi \circ (\mu_i \otimes 1))(m_{ij} \otimes n_j) \\
 &= \sum_{j=1}^k \phi(\mu_i(m_{ij}) \otimes n_j) \\
 &= \sum_{j=1}^k g(\mu_i(m_{ij}), n_j) \\
 &= \sum_{j=1}^k \mu'_i(g_i(m_{ij}, n_j)) \\
 &= \sum_{j=1}^k \mu'_i(m_{ij} \otimes n_j) \\
 &= p.
 \end{aligned}$$

As a result,  $\varinjlim (M_i \otimes N) \cong (\varinjlim M_i) \otimes N$ . □

**2.21.** Let  $(A_i)_{i \in I}$  be a family of rings indexed by a directed set  $I$ , and for each pair  $i \leq j$  in  $I$  let  $\alpha_{ij} : A_i \rightarrow A_j$  be a ring homomorphism, satisfying conditions (1) and (2) of Exercise 14. Regarding each  $A_i$  as a  $\mathbf{Z}$ -module we can then form the direct limit  $A = \varinjlim A_i$ . Show that  $A$  inherits a ring structure from the  $A_i$  so that the mappings  $A_i \rightarrow A$  are ring homomorphisms. The ring  $A$  is the direct limit of the system  $(A_i, \alpha_{ij})$ .

If  $A = 0$  prove that  $A_i = 0$  for some  $i \in I$ .

**Solution.** We define multiplication of  $A$  as follows. For any  $a, b \in A$ , by Exercise 2.15, there are some  $i \in I$  and some  $x_i, y_i \in A_i$  such that  $a = \alpha_i(x_i)$  and  $b = \alpha_i(y_i)$ . (We can say  $x_i, y_i$  lie on same  $A_i$  since  $I$  is a directed set; precisely, if  $x_{i_1} \in A_{i_1}$  and  $y_{i_2} \in A_{i_2}$ , then there exists  $i \in I$  such that  $i_1 \leq i$  and  $i_2 \leq i$ , and let  $x_i$  and  $y_i$  be  $\alpha_{i_1 i}(x_{i_1})$  and  $\alpha_{i_2 i}(y_{i_2})$ , respectively) Then define  $ab$  as  $\alpha_i(x_i y_i)$ . To show it is well-defined, suppose  $a = \alpha_j(x_j)$  and  $b = \alpha_j(y_j)$  for some  $j \in I$  and some  $x_j, y_j \in A_j$ . There exists some  $k \in I$  such that  $i \leq k$  and  $j \leq k$ , so

$$\alpha_k(\alpha_{ik}(x_i) - \alpha_{jk}(x_j)) = 0 \quad \text{and} \quad \alpha_k(\alpha_{ik}(y_i) - \alpha_{jk}(y_j)) = 0.$$

Then by Exercise 2.15, there is some  $k' \geq k$  so that<sup>4</sup>

$$\alpha_{kk'}(\alpha_{ik}(x_i) - \alpha_{jk}(x_j)) = 0 \quad \text{and} \quad \alpha_{kk'}(\alpha_{ik}(y_i) - \alpha_{jk}(y_j)) = 0.$$

Observe

$$\alpha_{ik}(x_i y_i) - \alpha_{jk}(x_j y_j) = \alpha_{ik}(x_i)(\alpha_{ik}(y_i) - \alpha_{jk}(y_j)) + \alpha_{jk}(y_j)(\alpha_{ik}(x_i) - \alpha_{jk}(x_j)).$$

Plugging it into the ring homomorphism  $\alpha_{kk'}$ , we obtain

$$\alpha_{kk'}(\alpha_{ik}(x_i y_i) - \alpha_{jk}(x_j y_j)) = 0.$$

This demonstrates that  $\alpha_i(x_i y_i) = \alpha_j(x_j y_j)$  in  $A$ , ensuring the well-defined nature of the multiplication. For  $i \in I$ , let  $1_i$  denote the multiplicative identity of  $A_i$ . As  $\alpha_{ij}(1_i) = 1_j$ , we can deduce that  $\alpha_i(1_i) = \alpha_j(1_j)$  for all  $i, j \in I$ . Let  $1$  represent  $\alpha_i(1_i)$ . Consequently, for any element  $a = \alpha_i(x_i)$  in  $A$ ,  $1a = \alpha_i(1_i x_i) = \alpha_i(x_i) = a$ . This confirms that the ring structure with which we have endowed  $A$  makes each  $\alpha_i$  a ring homomorphism.

Now suppose  $A = 0$ . Then for any  $i \in I$  and  $a_i \in A_i$ ,  $a_i$  is in  $\text{Ker}(\mu)$ , where  $\mu : C := \bigoplus_{i \in I} M_i \rightarrow M$  is the projection. Then as the solution of Exercise 2.15, there exists some  $j \in I$  such that  $\mu_{ij}(1_i) = 0$ . Since  $\mu_{ij}$  must send  $1_i$  to  $1_j$ , it implies  $A_j = 0$ .  $\square$

**2.22.** Let  $(A_i, \alpha_{ij})$  be a direct system of rings and let  $\mathfrak{N}_i$  be the nilradical of  $A_i$ . Show that  $\varinjlim \mathfrak{N}_i$  is the nilradical of  $\varinjlim A_i$ .

If each  $A_i$  is an integral domain, then  $\varinjlim A_i$  is an integral domain.

**Solution.** Let  $\mathfrak{N}$  denote the nilradical of  $\varinjlim A_i$ . Since  $\alpha_{ij}(\mathfrak{N}_i) \subseteq \mathfrak{N}_j$  for each  $i \leq j$ , the inclusion  $\iota_i : \mathfrak{N}_i \hookrightarrow A_i$  induces the corresponding homomorphism  $\iota : \varinjlim \mathfrak{N}_i \rightarrow \varinjlim A_i$ , and  $\iota$  is injective by Exercise 2.19. Therefore we can regard  $\varinjlim \mathfrak{N}_i$  as a subset of  $\varinjlim A_i$  via  $\iota$ . If  $x_i \in A_i$  is nilpotent, then  $\alpha_i(x_i)$  is also nilpotent, so  $\varinjlim \mathfrak{N}_i \subseteq \mathfrak{N}$  by Exercise 2.15. Conversely, suppose  $x \in \varinjlim A_i$  is nilpotent; i.e.,  $x^r = 0$  for some  $r \in \mathbb{Z}_{\geq 0}$ . By Exercise 2.15 again, it implies there exists some  $i, j$  such that  $x = \alpha_i(x_i)$  and  $\alpha_{ij}(x_i)^r = 0$ . Then  $\alpha_{ij}(x_i)$  is in  $\mathfrak{N}_j$ , so  $x = \alpha_j(\alpha_{ij}(x_i))$  is in  $\varinjlim \mathfrak{N}_i$ .  $\square$

**2.23.** Let  $(B_\lambda)_{\lambda \in \Lambda}$  be a family of  $A$ -algebras. For each finite subset  $J$  of  $\Lambda$  let  $B_J$  denote the tensor product (over  $A$ ) of the  $B_\lambda$  for  $\lambda \in J$ . If  $J'$  is another finite subset of  $\Lambda$  and  $J \subseteq J'$ , there is a canonical  $A$ -algebra homomorphism  $B_J \rightarrow B_{J'}$ . Let  $B$  denote the direct limit of the rings  $B_J$  as  $J$  runs through all finite subsets of  $\Lambda$ . The ring  $B$  has a natural  $A$ -algebra structure for which the homomorphisms  $B_J \rightarrow B$  are  $A$ -algebra homomorphisms. The  $A$ -algebra  $B$  is the *tensor product* of the family  $(B_\lambda)_{\lambda \in \Lambda}$ .<sup>5</sup>

**Solution.** There is nothing to do.  $\square$

**2.24.** If  $M$  is an  $A$ -module, the following are equivalent:

- i)  $M$  is flat;
- ii)  $\text{Tor}_n^A(M, N) = 0$  for all  $n > 0$  and all  $A$ -modules  $N$ ;
- iii)  $\text{Tor}_1^A(M, N) = 0$  for all  $A$ -modules  $N$ .

<sup>4</sup>Rigorously speaking, to show  $\alpha_{kk'}$  sends ‘both’ of them 0, we should repeat the same argument as showing that we can assume  $x_i$  and  $y_i$  lie on the same  $A_i$ .

**Remark.** An  $A$ -module  $P$  is **projective** if and only if for every surjective  $A$ -module homomorphism  $p : M \rightarrow M''$  and any  $A$ -module homomorphism  $h : P \rightarrow M''$ , there exists a lifting  $g$ ; that is, there exists a homomorphism  $g$  makes the following diagram commute:

$$\begin{array}{ccc} & P & \\ g \swarrow & \downarrow h & \\ M & \xrightarrow{p} & M'' \longrightarrow 0 \end{array}$$

It is easy to see that an  $A$ -module  $P$  is projective if and only if  $\text{Hom}_A(P, -)$  is an exact functor; that is, for every exact sequence of  $A$ -modules

$$0 \rightarrow M'' \rightarrow M \rightarrow M' \rightarrow 0,$$

the sequence

$$0 \rightarrow \text{Hom}(P, M'') \rightarrow \text{Hom}(P, M) \rightarrow \text{Hom}(P, M') \rightarrow 0$$

is also exact. For instance, every free  $A$ -module is projective ([3], Theorem 3.5).

For an  $A$ -module  $N$ , a **projective resolution** of  $N$  is an exact sequence

$$\cdots \rightarrow P_2 \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\varepsilon} N \rightarrow 0$$

in which each  $P_n$  is projective. If  $P_n$  is free, then the sequence is called **free resolution** of  $N$ . It is well known that every  $A$ -module  $N$  has a free resolution (the proof is actually not difficult, see Proposition 6.2 of [3]); hence, every  $A$ -module has a projective resolution. For a given projective resolution of  $N$ , remove  $N$

$$\cdots \rightarrow P_2 \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\partial_0} 0$$

and form the following sequence by tensoring it with  $M$ :

$$\cdots \rightarrow M \otimes_A P_2 \xrightarrow{1_M \otimes \partial_2} M \otimes_A P_1 \xrightarrow{1_M \otimes \partial_1} M \otimes_A P_0 \xrightarrow{1_M \otimes \partial_0} 0.$$

It is not an exact sequence in general, but it is easy to see that  $\text{Im}(1_M \otimes \partial_{n+1}) \subseteq \text{Ker}(1_M \otimes \partial_n)$  for all  $n \geq 0$ . Such sequence is called a **chain complex**. For  $n \geq 0$ , the  $A$ -module  $\text{Tor}_n^A(M, N)$  is the **homology** of this complex at position  $n$ ; that is,  $\text{Tor}_n^A(M, N) = \text{Ker}(1_M \otimes \partial_n) / \text{Im}(1_M \otimes \partial_{n+1})$  for  $n > 0$ , and  $\text{Tor}_0^A(M, N) = \text{Coker}(1_M \otimes \partial_1) \cong M \otimes_A N$ . Surprisingly,  $\text{Tor}_n^A(M, N)$  does not depend on the choice of projective resolution of  $N$  ([3], Proposition 6.20).

One of the most fundamental properties (in some context it is treated as an axiom for derived functors, which is the general notion of Tor functor; see Definition 2.1.1 of [4]) of Tor functor is as follows. If  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  is an exact sequence of  $A$ -modules, then for an  $A$ -module  $M$  there is a long exact sequence, called Tor exact sequence ([3], Theorem 6.27),

$$\begin{aligned} \cdots \rightarrow \text{Tor}_n^A(M, N') \rightarrow \text{Tor}_n^A(M, N) \rightarrow \text{Tor}_n^A(M, N'') \rightarrow \\ \text{Tor}_{n-1}^A(M, N') \rightarrow \text{Tor}_{n-1}^A(M, N) \rightarrow \text{Tor}_{n-1}^A(M, N'') \rightarrow \cdots \end{aligned}$$

which ends with

$$\cdots \rightarrow \text{Tor}_0^A(M, N') \rightarrow \text{Tor}_0^A(M, N) \rightarrow \text{Tor}_0^A(M, N'') \rightarrow 0.$$

Recall  $\text{Tor}_0^A(M, N) \cong M \otimes_A N$ .

**Solution.** [(i)  $\Rightarrow$  (ii)] Suppose an exact sequence

$$\cdots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow N \rightarrow 0$$

is a free resolution of  $N$ , and by tensoring with  $M$  we get

$$\cdots \rightarrow F_2 \otimes M \rightarrow F_1 \otimes M \rightarrow F_0 \otimes M \rightarrow N \otimes M \rightarrow 0.$$

Since  $M$  is flat, the resulting sequence is exact and therefore its homology groups, which are the  $\text{Tor}_n^A(M, N)$ , are zero for  $n > 0$ .

[(ii)  $\Rightarrow$  (iii)] It is trivial.

[(iii)  $\Rightarrow$  (i)] Let  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  be an exact sequence. Then from the Tor exact sequence,

$$\text{Tor}_1^A(M, N'') \rightarrow M \otimes N' \rightarrow M \otimes N \rightarrow M \otimes N'' \rightarrow 0$$

is exact. Since  $\text{Tor}_1^A(M, N'') = 0$  it follows that  $M$  is flat.  $\square$

**2.25.** Let  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  be an exact sequence, with  $N''$  flat. Then  $N'$  is flat  $\Leftrightarrow N$  is flat.

**Solution.** From the Tor exact sequence, we get an exact sequence

$$\cdots \rightarrow \text{Tor}_2(M, N'') \rightarrow \text{Tor}_1(M, N') \rightarrow \text{Tor}_1(M, N) \rightarrow \text{Tor}_1(M, N'') \rightarrow \cdots$$

for all  $A$ -modules  $M$ . If  $N''$  and  $N'$  are flat, then  $0 \rightarrow \text{Tor}_1(M, N) \rightarrow 0$  is exact, implying  $\text{Tor}_1(M, N) = 0$ . Therefore  $N$  is flat. If  $N''$  and  $N$  are flat, then  $0 \rightarrow \text{Tor}_1(M, N') \rightarrow 0$  is exact. As a result  $N'$  is flat.  $\square$

**2.26.** Let  $N$  be an  $A$ -module. Then  $N$  is flat  $\Leftrightarrow \text{Tor}_1(A/\alpha, N) = 0$  for all finitely generated ideals  $\alpha$  in  $A$ .

**Solution.** If  $N$  is flat, then  $\text{Tor}_1(M, N) = 0$  for all  $A$ -modules  $M$  by Exercise 2.24. To show the converse, firstly we claim that  $N$  is flat if  $\text{Tor}_1(M, N) = 0$  for all finitely generated  $A$ -modules  $M$ . Let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence of finitely generated  $A$ -modules. Then from the Tor exact sequence, we get an exact sequence

$$\text{Tor}_1(M'', N) \rightarrow M' \otimes N \rightarrow M \otimes N \rightarrow M'' \otimes N \rightarrow 0.$$

Since  $\text{Tor}_1(M'', N) = 0$  by the assumption, we conclude that for any injective homomorphism  $f : M' \rightarrow M$  the corresponding homomorphism  $f \otimes 1 : M' \otimes N \rightarrow M \otimes N$  is injective. Hence  $N$  is flat by Proposition 2.19, and this shows the claim holds. Now suppose  $\text{Tor}_1(A/\alpha, N) = 0$  for all finitely generated ideals  $\alpha$  in  $A$ . If  $M$  is finitely generated, let  $x_1, \dots, x_n$  be a set of generators of  $M$ , and let  $M_i$  be the submodule generated by  $x_1, \dots, x_i$ . Observe that for a given cyclic module  $Ax$ , a map  $f : A \rightarrow Ax$  given by  $1 \mapsto x$  is an  $A$ -module homomorphism, implying  $Ax \cong A/\text{Ker}(f)$ . Since  $M_i/M_{i-1}$  is generated by a single element for  $2 \leq i \leq n$ ,  $M_i/M_{i-1} \cong A/\alpha_i$  for some ideal  $\alpha_i$ . Consider the exact sequence

$$0 \rightarrow M_{i-1} \rightarrow M_i \rightarrow M_i/M_{i-1} \rightarrow 0$$

for  $2 \leq i \leq n$ . Since  $\text{Tor}_1(M_1, N) = 0$  and  $\text{Tor}_1(M_i/M_{i-1}, N) = 0$  by the hypothesis for  $2 \leq i \leq n$ , proceeding by induction on  $i$  we get  $\text{Tor}_1(M, N) = \text{Tor}_1(M_n, N) = 0$ . This ends the proof.  $\square$

**2.27.** A ring  $A$  is *absolutely flat* if every  $A$ -module is flat. Prove that the following are equivalent:

- i)  $A$  is absolutely flat
- ii) Every principal ideal is idempotent.
- iii) Every finitely generated ideal is a direct summand of  $A$ .

**Solution.** [i]  $\Rightarrow$  ii] Let  $x \in A$ . Since  $A/(x)$  is a flat  $A$ -module, the map  $\alpha : (x) \otimes A/(x) \rightarrow A \otimes A/(x) = A/(x)$  induced by the inclusion  $(x) \hookrightarrow A$  is injective. Since  $\alpha(x \otimes \bar{a}) = x\bar{a} = 0$ , we get  $(x) \otimes A/(x) = 0$ . However,  $(x) \otimes A/(x) = (x)/(x^2)$  by Exercise 2.2, so  $(x) = (x^2)$ .

[ii]  $\Rightarrow$  iii] Let  $x \in A$ . Then  $x = ax^2$  for some  $a \in A$ , hence  $e = ax$  is idempotent, and  $(x) = (e)$  because  $x = xe$ . Now if  $e, f$  are idempotents, then  $(e, f) = (e + f - ef)$  since  $e(e + f - ef) = e$  and  $f(e + f - ef) = f$ . Therefore every finitely generated ideal is principal, and generated by an idempotent  $e$ , hence is a direct summand because  $A = (e) \oplus (1 - e)$ .<sup>5</sup>

[iii]  $\Rightarrow$  i] Clearly  $A$  is an flat  $A$ -module, so every finitely generated ideal of  $A$  is flat by Exercise 2.4. Since  $A/\alpha$  is a direct summand of  $A$  for any finitely generated ideal  $\alpha$ , we have  $\text{Tor}_1(A/\alpha, N) = 0$  for any  $A$ -module  $N$ . By Exercise 2.26, every  $A$ -module is flat.  $\square$

**2.28.** A Boolean ring is absolutely flat. The ring of Chapter 1, Exercise 7 is absolutely flat. Every homomorphic image of an absolutely flat ring is absolutely flat. If a local ring is absolutely flat, then it is a field.

If  $A$  is absolutely flat, every non-unit in  $A$  is a zero-divisor.

**Solution.** Every principal ideal of a Boolean ring is clearly idempotent, which makes it absolutely flat.

Let  $A$  be the ring of Chapter 1, Exercise 7; i.e.,  $A$  is a nonzero ring in which every element  $x$  satisfies  $x^n = x$  for some  $n > 1$  (depending on  $x$ ). Suppose  $x^n = x$  and  $y^m = y$  for some  $n, m > 1$ . Then  $x = x(x^{n-1} + y^{m-1} - x^{m-1}y^{m-1})$  and  $y = y(x^{n-1} + y^{m-1} - x^{m-1}y^{m-1})$ , so  $(x, y) = (x^{n-1} + y^{m-1} - x^{m-1}y^{m-1})$ . Therefore every finitely generated ideal is principal. Since  $x = x^{n-2}x^2$ , we have  $(x) = (x^2)$ , so  $A$  is absolutely flat.

Let  $\phi : A \rightarrow B$  be a ring homomorphism where  $A$  is absolutely flat. Then  $\phi(A) \cong A/\text{Ker}(\phi)$ . For any principal ideal  $(\bar{x})$  of  $A/\text{Ker}(\phi)$ , clearly  $(\bar{x}) = (\bar{x})^2$  because  $(x) = (x)^2$  in  $A$ . Therefore every homomorphic image of an absolutely flat ring is absolutely flat.

Suppose  $A$  is a local ring which is absolutely flat. For any  $x \in A$ , since every principal ideal is idempotent, we have  $x = ax^2$  for some  $a \in A$ . Then  $e = ax$  is idempotent, but a local ring contains no idempotent neither 0 nor 1. Therefore, if  $x$  is nonzero,  $x$  is a unit, so  $A$  is a field.

Now suppose  $A$  is absolutely flat and  $x$  is a non-unit in  $A$ . Since every principal ideal is idempotent, there is some  $a \in A$  so that  $x(1 - ax) = x - ax^2 = 0$ . If  $x$  is not a zero divisor, then  $1 - ax = 0$ , leading to a contradiction. This shows that every non-unit in  $A$  is a zero-divisor.  $\square$

<sup>5</sup>Consider maps  $A \rightarrow (e) \oplus (1 - e)$  and  $(e) \oplus (1 - e) \rightarrow A$  given by  $a \mapsto (ae, a(1 - e))$  and  $(a, b) \mapsto a + b$ , respectively. Then they are two-sided inverses of each other.

## References

- [1] K. Conrad. Exterior powers. <https://kconrad.math.uconn.edu/blurbs/linmultialg/extmod.pdf>.
- [2] Balazs Strenner ([https://mathoverflow.net/users/11214/balazs\\_strenner](https://mathoverflow.net/users/11214/balazs_strenner)). Atiyah-macdonald, exercise 2.11. <https://mathoverflow.net/q/47846>.
- [3] Joseph J. Rotman. *An Introduction to Homological Algebra*. Springer Science+Business Media, second edition, 2009.
- [4] Charles A. Weibel. *An Introduction to Homological Algebra*. Cambridge University Press, 1994.